



Green Tea Extract Reduced Lipopolysaccharide-Induced Inflammation in L2 Cells as Acute Respiratory Distress Syndrome Model Through Genes and Cytokine Pro-Inflammatory

Didik Priyandoko ¹, Wahyu Widowati ^{2*}, Lenny Lenny ³, Sintya Novianti ³, Revika Revika ³, Hanna Sari Widya Kusuma ⁴ and Ika Adhani Sholihah ^{4,5}

1. Biology Study Program, Faculty of Mathematics and Natural Sciences, Indonesia University of Education, Bandung 40154, Indonesia

2. Faculty of Medicine, Maranatha Christian University, Bandung 40164, Indonesia

3. Faculty of Biotechnology, Atma Jaya Catholic University of Indonesia, BSD Campus, Tangerang 15345, Indonesia

4. Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung 40163, Indonesia

5. School of Life Sciences and Technology, Bandung Institute of Technology, Bandung 40132, Indonesia

Abstract

Background: Acute Respiratory Distress Syndrome (ARDS) is a severe lung inflammatory condition that has the capacity to impair gas exchange and lead to hypoxemia. This condition is found to have been one of the most prevalent in patients of COVID-19 with a more serious condition. Green tea (*Camellia sinensis* L.) contains polyphenols that possess many health benefits. The purpose of this study was to assess the anti-inflammatory activities of green tea extract in Lipopolysaccharide (LPS)-induced lung cells as ARDS cells model.

Methods: In this study, rat lung cells (L2) were induced by LPS to mimic the inflammation observed in ARDS and later treated with green tea extract. Pro-inflammatory cytokines such as Interleukin (IL)-12, C-Reactive Protein (CRP) as well as Tumor Necrosis Factor- α (TNF- α) were investigated using the ELISA method. Gene expression of NOD-Like Receptor Protein 3 (*NLRP-3*), Receptor for Advanced Glycation End-product (RAGE), Toll-like Receptor-4 (*TLR-4*), and Nuclear Factor-kappa B (*NF- κ B*) were evaluated by qRT-PCR. Apoptotic cells were measured using flow cytometry.

Results: The results showed that green tea extract treatment can reduce inflammation by suppressing gene expressions of *NF- κ B*, *NLRP-3*, *TLR-4*, and *RAGE*, as well as pro-inflammatory cytokines such as IL-12, TNF- α , and CRP, an acute phase protein. Apoptosis levels of inflamed cells also found to be lowered when green tea extract was administered; thus, also increasing live cells compared to non-treated cells.

Conclusion: These findings could lead to the future development of supplements from green tea to help alleviate ARDS symptoms, especially during critical moments such as the current pandemic.

Keyword: Acute respiratory distress syndrome, *Camellia sinensis*, Cytokines, Inflammation, Tea

To cite this article: Priyandoko D, Widowati W, Lenny L, Novianti S, Revika R, Kusuma HSW, et al. Green Tea Extract Reduced Lipopolysaccharide-Induced Inflammation in L2 Cells as ARDS Model Through Genes and Cytokine Pro-Inflammatory. Avicenna J Med Biotech 2024;16(1):57-65.

* Corresponding author:
Wahyu Widowati, Ph.D., Faculty
of Medicine, Maranatha Christian
University, Bandung 40164,
Indonesia
Tel: +62 819 10040010
E-mail:
wahyu_w60@yahoo.com
Received: 9 May 2023
Accepted: 23 Aug 2023

Introduction

Systemic inflammation of the lungs of the body causes Acute Respiratory Distress Syndrome (ARDS). ARDS is often associated with diseases such as traumatic brain injury, burn injury, immune system systemic suppression, and bacterial or viral infections, as recently found out in the case of SARS-CoV-2 ¹⁻³. ARDS is characterized by very poor oxygenation, reduced pulmonary infiltrates, and bilateral radiographic

infiltrates ⁴. Various studies have shown that the death of patients with ARDS is common ⁵. Previous studies have shown that there are similarities between the pathological anatomy of patients who died from COVID-19 and ARDS. The pathological anatomy of ARDS can aid in the understanding and treatment of COVID-19 patients ⁶.

ARDS is usually caused and worsened by a phe-

nomenon known as the Cytokine Storm Syndrome (CS). CS is a phenomenon in which the immune cells, as a direct response to injury or pathogens, release a huge amount of pro-inflammatory cytokine and chemokine like Interleukin (IL)-6, IL-12, IL-1 β , and Tumor Necrosis Factor- α (TNF- α) in an attempt to attract other immune cells to migrate to the site of the injury^{5,7,8}. Moreover, C-Reactive Protein (CRP), which is also released in reaction to inflammation and primarily functions to aid in opsonization as well as the release of other pro-inflammatory cytokines, is also produced⁹. The genes that are responsible for producing and playing a role in its cascade in the inflammatory pathways are also diverse. These inflammatory cytokines would then be recognized by receptors which are regulated by genes such as Toll-Like Receptor-4 (TLR-4) and Receptor for Advanced Glycation End products (RAGE)¹⁰. The gene that is also responsible for the production of cytokines would include Nod-Like Receptors-3 (NLRP-3) which also plays a role in the pathway of "Nuclear Factor-kappa B (NF- κ B)"^{11,12}. Accordingly immune cells, whether innate or adaptive, will migrate and destroy the inflammation trigger and protect infected or injured cells^{13,14}. At an appropriate amount, these immune cells will help to cure, but when there is too much destruction our body will not be able to handle it. Especially in the case of ARDS where the destruction happens in one of the most critical tissues in the body, the alveoli which help with gas exchange. Once the epithelial cells are destroyed, leakage of high-protein fluids will lead to edema. Over time, this edema can cause hypoxia due to its impact on gas exchange^{1,3}.

Treatment for patients usually include the usage of low tidal mechanical ventilators in hospitals. However, other than synthetic drugs such as simvastatin, and in the case of COVID-19, methylprednisolone and dexamethasone, an alternative cure has yet to be found^{15,16}. The problem with synthetic drugs is that they often cause negative side consequences. Non-Steroidal. Anti-Inflammatory Drugs (NSAIDs), generally also used to cure inflammations, are known to increase the risk of gastrointestinal illnesses like dyspepsia and esophagitis^{17,18}. Various plants and herbs have been investigated as the source of anti-inflammatory agents. Tea (*Camellia sinensis* L.) is a renowned plant that contains large amounts of polyphenols and can act as an anti-inflammatory agent. Tea contains several types of compound groups such as phenolics, alkaloids, polysaccharides, and saponins¹⁹. The main polyphenol compound in green tea that is thought to help with its anti-inflammation activities are flavonoids such as quercetin and myricetin. A previous study has also detected 8-prenylnaringenin (8-PN) and 6-prenylnaringenin (6-PN) in the Green Tea Extract (GTE) used^{17,20}.

The aim of this study was to evaluate the effect of GTE on Lipopolysaccharide (LPS)-induced lung cells as ARDS cells model by measuring gene expression of

NF- κ B, NLRP-3, RAGE, and TLR-4 by RT-PCR and examination of proinflammatory cytokine such as cytokines TNF- α , CRP, and IL-12 using sandwich Enzyme-Linked Immunosorbent Assay (ELISA). Flow cytometry was utilized to examine the apoptotic activity of the cells to determine their condition.

Materials and Methods

Green tea extract preparation

The GTE was processed by PT. Fathonah Amanah Shidiq Tabligh (Depok, Indonesia) (Batch No. 00107201057) and the process based on Good Manufacturing Practice (GMP) standard by The Indonesian Food and Drug Authority (BPOM). The extraction process of GTE used 70% ethanol as its solvent. Before GTE was stored at 25 $\pm$ 2 °C, lactose was added to GTE first for getting the powder form.

L2 cells-LPS induced as ARDS cells model

Rat lung (L2) cells (ATCC® CCL-149™) was obtained from the Biomolecular and Biomedical Research Center, Aretha Medika Utama which is located in Bandung, West Java, Indonesia. T75 flask with a complete medium was used for cell growth until confluent. The complete medium composition was Dulbecco's Modified Eagle's Medium (DMEM) High-glucose (Biowest, L1003-500), 10% Fetal Bovine Serum (FBS) (Biowest, S1810-5000), 1% Antibiotic-antimycotic (ABAM) 100 $\times$ (Biowest, L0009), and 0.1% Gentamicin (Gibco, 15750078). Confluent cells were induced with 4 μ g/ml LPS *Escherichia coli* O55:B5 (Sigma-Aldrich L2880) to produce an inflammation model and incubated for 18 hr at 37°C in a 5% CO₂ incubator. Various concentrations of GTE (1.56, 3.13, and 6.25 μ g/ml) were applied to the cells and incubated for 24 hr in a 5% CO₂ incubator (37°C). The levels of IL-6, CRP and TNF- α in the harvested supernatant cells were measured using the ELISA method, while the cell pellets were collected for RNA isolation to be used for testing TLR-4, RAGE, NF- κ B, and NLRP-3 gene expression using qRT-PCR⁸.

Quantification of TNF- α , CRP, and IL-12 levels in the L2 cells using ELISA

The supernatant cells were used to determine the levels of pro-inflammatory cytokine expression in the L2 cells, such as TNF- α , CRP, and IL-12. The TNF- α , CRP, and IL-12 levels were determined by the ELISA kit (Elabscience E-EL-R0019, E-EL-R0506, and E-EL-R0064, respectively). The process was carried out according to the manufacturer's guidelines. The sample absorbances were tested at a wavelength of 450 nm⁸.

Quantification of NF- κ B, NLRP-3, RAGE, and TLR-4 gene expression in the L2 cells using qRT-PCR

The levels of NF- κ B, NLRP-3, RAGE, and TLR-4 gene expression were determined from pellet cells. Direct-zol RNA Miniprep Plus Kit (Zymo, R2073) was applied to isolate total RNA, the procedure was according to the protocol of manufacture. Then, the Sensi-

FAST cDNA synthesis kit was used to synthesize complementary DNA and the procedure was according to the protocol of manufacture. AriaMx 3000 Real-Time PCR System (Agilent) was used to measure quantitative gene expression, primer sequence (Table 1), and Evagreen master mix (Bio-Rad, 1725200) for the reaction mixture. The procedure was undertaken based on the manufacturer's protocol ²¹. Real-time PCR was run for 40 cycles using *GAPDH* as reference gene, and the annealing temperatures was 54°C, 56°C, and 57°C for *TLR-4*, *RAGE*, and *NF-κB*, *NLRP-3*, respectively.

Apoptosis assay using flow cytometry analysis

Annexin V-FITC/PI Apoptosis Detection Kit (Elabsci, E-CK-A211) was utilized to assess the cells apoptosis in line with the manufacturer's instructions, with slight modification. Cells incubated for 2 hr on a 6-well plate (n=500,000). After 4 days, the growth medium was removed, harvested and centrifuged at 1,600 rpm for 5 min, and then 500 µl FACS buffer added on pellet and centrifuged. Additionally, annexin and Propidium Iodide (PI) staining was performed on the pellets after the addition of 100 µl FACS buffer. MACS-Quant Analyzer. 10 Flow Cytometer (Miltenyi, Bergisch Gladbach, Germany) was used to evaluate the stained cells that have been incubated for 1 hr ²².

Statistical analysis

The data analysis was conducted using SPSS 26.0 software, and experimental results are shown as mean ± SD. The comparison between groups was assessed using a one-way ANOVA and the post hoc test was Dunnett's test and Tukey HSD test with p<0.05. Analysis using Kruskal-Wallis and Mann-Whitney. post hoc with p<0.05 was used for non-parametric.

Results

The influence of GTE on *TNF-α*, *CRP*, and *IL-12* levels

Pro-inflammatory cytokine was increased by LPS induction. Figure 1 shows the effect of GTE toward IL-12, *TNF-α*, and *CRP* levels. Compared to the negative control, the level of IL-12, *TNF-α*, and *CRP* increased significantly in positive control (p<0.05). Treatment




Figure 1. Effect of GTE toward IL-12, *TNF-α*, *CRP* on LPS-induced L2 cells. A) IL-12 level (pg/ml) on LPS-induced L2 cells, B) *TNF-α* level (pg/ml) on LPS-induced L2 cells, C) *CRP* level (ng/ml) on LPS-induced L2 cells.

* The data is presented as means value ± standard deviation. NC: untreated cell, PC: LPS-induced cell, DC: PC+DMSO1%, GTE 6.25: PC + GTE 6.25 µg/ml, GTE 3.13: PC + GTE 3.13 µg/ml, GTE 1.65: PC + GTE 1.65 µg/ml. A different mark (alphabetical) indicates a significant difference among treatments toward *CRP* level based on Kruskal-Wallis and Mann-Whitney post hoc test (p<0.05). A different letter (alphabetical) indicates a significant difference among treatments based on Anova and Tukey HSD post hoc test for *TNF-α* level and Dunnett T3 post hoc for IL-12 level at Anova and Tukey HSD post hoc test p<0.05.

with GTE on L2 cells reduced pro-inflammatory cytokine levels compared to positive control. The results showed that GTE reduced *CRP* and *TNF-α* levels significantly in the rat lung cell at concentration of 1.56 µg/ml (p<0.05). Meanwhile, the IL-12 level did not show any significant reduction in all concentration of GTE.

Effect of GTE toward *TLR-4*, *RAGE*, *NF-κB*, and *NLRP-3* genes expression

LPS-induced L2 cells significantly increased *TLR-4*, *RAGE*, *NF-κB*, and *NLRP-3* genes expression (p<0.05).

Table 1. Primer sequence

Gene	Primer sequence (5'-3')	Product length (bp)	Reference
	Forward/Reverse		
<i>TLR-4</i>	AGTGAGAATGCTAAGGTTGG	289	NCBI Reference Sequence: NM_019178.2
	ATTAGGAAGTACCTCTATGCAG		
<i>RAGE</i>	CGAGTCTACCAGATTCTCTG	163	NCBI Reference Sequence: NM_053336.2
	CTTTGCCATCAGGAATCAG		
<i>NF-κB</i>	GGACTATGACTTGAATGCGG	230	NCBI Reference Sequence: NM_199267.2
	ACACCTCAATGCTCTTCTTCTG		
<i>NLRP-3</i>	CAGATGCTGGAGTTAGACAACCTG	153	NCBI Reference Sequence: NM_001191642.1
	TTCAGAACCTCACAGAGCGT		
<i>GAPDH</i>	TCAAGATGGTGAAGCAG	217	NCBI Reference Sequence: NM_001289726.1
	ATGTAGGCCATGAGGTCCAC		

As shown in figure 2, GTE treatments downregulated the expression of *TLR-4*, *RAGE*, *NF- κ B*, and *NLRP-3* genes expression ($p < 0.05$). The results showed that *TLR-4*, *RAGE*, and *NLRP-3* genes expression were significantly decreased with GTE treatment at the concentration of 1.56 $\mu\text{g/ml}$ ($p < 0.05$). In contrast, the 6.25 $\mu\text{g/ml}$ GTE showed the most effective concentration in suppressing *NF- κ B* gene expression but did not show any significant difference with the positive control.

Effect of GTE toward cells apoptosis

The effect of GTE was analyzed by flow cytometry in triplicate using concentrations of 6.25, 3.13, and 1.56 $\mu\text{g/ml}$ respectively. Figure 3 shows that LPS induction increased the number of cells in necrosis, early, and late apoptosis, and also reduces the number of live cells quantity compared to the negative control. The dot blot of the effect of GTE on cell apoptosis can be seen in figure 4. The number of necrosis, early and late apoptosis were decreased following GTE treatment, while the number of viable cells increased.

According to figure 4, the GTE treatment increased the percentage of live cells compared to the positive control. Moreover, depending on the concentration, treatments with GTE decreased the percentage of necrosis cells with concentration dependent manner, and GTE treatment reduced the percentage of apoptotic cells. The highest GTE concentration (6.25 $\mu\text{g/ml}$) had the smallest apoptosis percentage compared to other GTE treatments with the apoptosis percentage of 10.34%.

Discussion

One of the life-threatening illnesses observed in severely ill COVID-19 patients is ARDS. ARDS, just




Figure 2. Effect of GTE toward *TLR-4*, *RAGE*, *NF- κ B*, and *NLRP-3* gene expression on LPS-induced L2 cells. A) *TLR-4* gene expression on LPS-induced L2 cells, B) *RAGE* gene expression on LPS-induced L2 cells, C) *NF- κ B* gene expression on LPS-induced L2 cells, D) *NLRP-3* gene expression on LPS-induced L2-cells.

* The data is presented as means value $\pm$ standard deviation. NC: untreated cell, PC: LPS-induced cell, DC: PC+DMSO 1%, GTE 6.25: PC + GTE 6.25 $\mu\text{g/ml}$, GTE 3.13: PC + GTE 3.13 $\mu\text{g/ml}$, GTE 1.65: PC + GTE 1.65 $\mu\text{g/ml}$. A different mark (alphabetical) indicates a significant difference among treatments toward *RAGE* gene expression, based on Kruskal-Wallis and Mann-Whitney post hoc test ($p < 0.05$). A different letter (alphabetical) indicates a significant difference among treatments based Anova and Tukey HSD post hoc test for *NLRP-3* gene expression and based Anova and Dunnett T3 post hoc test for *TLR-4* and *NF- κ B* gene expression ($p < 0.05$).

like any other inflammation can be induced by different agents such as protein in viruses or LPS of bacteria.




Figure 3. Effect of GTE toward early apoptosis, live cells, necrosis cells, late apoptosis on LPS-induced L2 cells.

* The data is presented as means value $\pm$ standard deviation. NC: untreated cell, PC: LPS-induced cell, DC: PC+DMSO 1%, GTE 6.25: PC + GTE 6.25 $\mu\text{g/ml}$, GTE 3.13: PC + GTE 3.13 $\mu\text{g/ml}$, GTE 1.65: PC + GTE 1.65 $\mu\text{g/ml}$. The number of early apoptosis, live, necrosis, and late apoptosis cells was measured in triplicate for each sample. A) Early apoptosis on LPS-induced L2 cells, B) Live cells on LPS-induced L2 cells, C) Necrosis cells on LPS-induced L2 cells, D) Late apoptosis on LPS-induced L2 cells. The data is presented as means value $\pm$ standard deviation. NC: untreated cell, PC: LPS-induced cell, DC: PC+DMSO 1%, GTE 6.25: PC + GTE 6.25 $\mu\text{g/ml}$, GTE 3.13: PC + GTE 3.13 $\mu\text{g/ml}$, GTE 1.65: PC + GTE 1.65 $\mu\text{g/ml}$. A different mark (alphabetical) indicates a significant difference among treatments toward early apoptosis, live cells, necrosis cells, late apoptosis based on Anova and Tukey HSD post hoc test ($p < 0.05$).




Figure 4. The representative dot blots of different GTE concentrations on LPS-induced L2 cells toward percentage of apoptosis by flow cytometry. * A) Negative control (untreated cell): live cells 90.23%, early apoptosis 3.47%, late apoptosis 1.03%, necrosis cells 5.28%. B) Positive control (LPS-treated cells): live cells 75.39%, early apoptosis 11.38%, late apoptosis 4.57%, necrosis cells 8.66%. C) DMSO control (untreated cells+DMSO): live cells 85.48%, early apoptosis 7.30%, late apoptosis 3.00%, necrosis cells 4.22%. D) GTE 1.56 $\mu\text{g/ml}$: live cells 80.55%, early apoptosis 12.82%, late apoptosis 3.06%, necrosis cells 3.57%. E) GTE 3.13 $\mu\text{g/ml}$: live cells 82.78%, early apoptosis 10.75%, late apoptosis 3.22%, necrosis cells 3.25%. F) GTE 6.25 $\mu\text{g/ml}$: live cells 82.48%, early apoptosis 10.45%, late apoptosis 3.27%, necrosis cells 3.08%.

LPS administration in both *in vivo* and *in vitro* leads to inflammation and causes the release of inflammatory cytokines^{23,24}. LPS induction led to an increase in CRP levels as well as pro-inflammatory cytokines as TNF- α and IL-12, as shown in figure 1. This is also proven by a previous study, who also found that LPS induction releases uncontrolled cytokines that could lead to lung injury²⁵.

In order to compare the cytokine or expression levels of inflamed cells with those of healthy cells, negative controls were required for all assays in which LPS or the extract was not added to the cells. Additionally, both positive and DMSO controls were established. To confirm that the solvent for the GTE had no significant effect on the inflamed cells, DMSO was used as a control. This would allow us to conclude that the difference in results between treated and untreated cells was due to the GTE treatment and not the solvent used.

In the absence of a stimulus, healthy cells should have lower levels of pro-inflammatory cytokines and expression of genes related to inflammation than inflamed cells²⁶. When pro-inflammatory triggers such as LPS are identified, pro-inflammatory cytokines begin to be produced. In both animal models and patient studies, the *TLR-4* has been shown to be a receptor for the response to LPS²⁷. *RAGE* receptors, receptors highly expressed in skin and lung tissue, are also

involved in the LPS response. *RAGE* expression can be increased in the presence of inflammatory conditions, aging, and at the time of injury. *RAGE* and *TLRs* are crucial components of the natural immune system; as these compounds can interact with various microbial products and molecules (one of them is LPS) in inflammation and tissue injury²⁸. The *NF- κ B* signaling pathway can be activated by either *TLR-4* or *RAGE* receptors. Activated *NF- κ B* controls the synthesis of proinflammatory cytokines and chemokines during inflammation²⁹. Previous research has shown that *NF- κ B* signaling is involved in activating *NLRP-3*³⁰. An immunological complex called the *NLRP-3* inflammasome is made up of *NLRP-3*, an apoptosis-associated protein with a pro-caspase-1 and a C-terminal caspase recruitment domain (ASC)³¹. *NLRP-3* also recognizes Reactive Oxygen Species (ROS) and Damage-Associated Molecular Patterns (*DAMPs*) in inflammation cells (LPS-induced cells)³².

Figures 1-4 show that DMSO controls had insignificant levels of pro-inflammatory cytokines compared to positive controls. Based on the results of DMSO control for the gene expression, levels of *TLR4*, *RAGE*, *NF- κ B*, and *NLRP3* were comparable with figure 3; where the DMSO control had insignificant results with positive control. Thus, it could be assumed that all the results from treatments are purely based on the

treatment of GTE.

Green tea contains many beneficial polyphenols such as catechins, quercetin, myricetin, and kaempferol³³. These polyphenols have made green tea a well-known beverage with many therapeutic benefits and are often used in many traditional therapies. Additionally, green tea has been reported to have antioxidant and anti-inflammatory properties that can help decrease inflammation by lowering inflammatory markers. Its antioxidant activity helps to scavenge ROS which can lead to the suppression of *NF-κB* activity^{34,35}. Figure 2 shows that there was an increase in the expression of *TLR-4*, *RAGE*, *NF-κB*, and *NLRP-3* genes. LPS induction on human dental pulp stem cells upregulated *TLR-4* gene expression³⁶. The findings of this study demonstrate that *TLR-4* is a particular receptor for inducing LPS in cells³⁷. In addition, in this study, the increase of *TLR-4* gene expression was in line with *NF-κB* gene expression in response to LPS. These results are consistent with previous studies which LPS induction increased *NF-κB* gene expression³⁷. Moreover, LPS induction also increases the *NLRP-3* expression on RAW 264.7 cells and is in line with the *NF-κB* expression³⁸. The results of this research indicate that GTE down-regulated the expression of *TLR-4*, *RAGE*, *NF-κB*, and *NLRP-3* mRNA levels in LPS-induced L2 cells as ARDS cells model.

The findings in figure 2 concur with those of the previous investigations. From figure 2, it can be inferred that GTE has the ability to suppress the mRNA expression of *TLR-4*, *RAGE*, *NF-κB*, and *NLRP-3* in the LPS-induced L2 cells as ARDS cells model. The expression of proinflammatory cytokines including *TNF-α* and *IL-12* was also decreased by reducing *NF-κB* and other gene expressions (Figure 2).

Besides *NF-κB*, green tea catechin also inhibits the pro-inflammatory mediators' secretion through *NLRP-3* inflammasome, a multiprotein complex that activates inflammatory responses³⁹. In prior research, treatment with quercetin, one of the components in green tea, was shown to be able to decrease both the mRNA level of *NLRP-3* and the amount of the pro-inflammatory cytokine⁴⁰. Epigallocatechin-3-gallates (EGCG), in particular, was discovered in a different study to have the ability to suppress and lower the expression of *TLR-4* and *RAGE* following a pre-treatment of EGCG in rat LPS-induced inflammation⁴¹⁻⁴³. Inhibiting those gene expressions also resulted in a reduction of pro-inflammatory cytokines *TNF-α* and *IL-12*^{43,44}.

Precisely in this study, GTE treatment at all three concentrations significantly reduced *TNF-α* levels as compared to positive controls. These matched previous findings of GTE and their ability to lower inflammation by modulating pro-inflammatory cytokines like *TNF-α*, is especially due to the presence of flavonoids such as myricetin and quercetin⁴⁵. Treatment with 1.56 $\mu\text{g/ml}$ GTE showed the most active to lower *TNF-α*

level compared to other GTE treatments and significantly decreased *TNF-α* level compared to positive control. Moreover, GTE 1.56 $\mu\text{g/ml}$ also decreased *IL-12* level significantly compared to positive control.

The CRP levels were also found to be highly associated to immunological alterations as a marked rise in pro-inflammatory biomarkers⁴⁶. Although CRP is mostly produced in the liver, its mRNA has been found in respiratory tract epithelial cells and T-lymphocytes⁴⁷. The cytokine storm in ARDS induces macrophage activation syndrome and results in a pro-inflammatory hypercytokinemia profile, which increases CRP generation by hepatocytes^{48,49}.

Based on the results shown in figure 2C, administration of GTE showed a significantly reduced level of CRP. The decreased level of CRP is in correlation with *IL-6* and *TNF-α* levels. CRP, an acute phase protein, was induced by both *IL-6* and *TNF-α* in inflammation. It was shown that by decreasing these inflammatory cytokines, the level of CRP was reduced⁹. As discussed previously, GTE reduced the level of inflammatory mediators in an inflammation cells model. The most effective result was seen in 1.56 $\mu\text{g/ml}$ GTE for all parameters, except for the expression of *NF-κB* in which 6.25 $\mu\text{g/ml}$ GTE was the most active to decrease *NF-κB* gene expression. Therefore, further research is required to establish the excellent concentration of GTE for its effectiveness in treating ARDS.

One other marker of inflammation is apoptosis or programmed cell death⁵⁰. Apoptosis is very crucial when it comes to the host defense system as it would allow the damaged or infected cells to die and end further damage inside the host⁵⁰. There are several ways apoptosis can be activated and in inflammation, apoptosis could happen due to both internal aspects such as mitochondrial dysfunction and stress, along with external factors including induction by LPS^{50,51}, both of which activate caspase to start the apoptosis⁵². This is why in most cases of inflammation the apoptosis level of cells is usually increased⁵¹.

In case of ARDS, the death of a large number of cells could also lead to the destruction of vital tissues, causing what would ideally be a preventative process of a deadly damage rapidly^{50,51,53}. Apoptosis suppression has played a crucial role in alleviating inflammation symptoms. The result of the apoptosis assay by flow cytometry indicated that all three-concentrations managed to decrease apoptosis levels of inflamed cells as could be seen in the significant difference of live cells compared to the positive controls. The results of the study were consistent with previous study⁵⁴ which found that quercetin, one of the constituents of green tea can suppress apoptosis in LPS-induced inflammation. Moreover, all GTE concentration treatments showed significantly reduced cell death, early and late apoptosis levels.

Conclusion

In this study, it is proven that GTE can help reduce the inflammation by modulation of TNF- α , CRP and IL-12 pro-inflammatory marker as well as suppressing inflammation-related genes expression including *TLR-4*, *RAGE*, *NF- κ B*, and *NLRP-3*. Furthermore, GTE was able to reduce the apoptosis in inflamed cells.

Acknowledgement

Financial assistance was provided by the Minister of Education, Culture, Research, and Technology of the Republic of Indonesia (Penelitian Dasar Unggulan Perguruan Tinggi 2021) under grant number 163/E4.1/AK.04.PT/2021. The Biomolecular and Biomedical Research Center, Aretha Medika Utama, Bandung, West Java, Indonesia, also provided the laboratory resources and research methodology for this study.

Conflict of Interest

The authors declare that they have no competing interest.

References

- Han S, Mallampalli RK. The acute respiratory distress syndrome: from mechanism to translation. *J Immunol* 2015;194(3):855-60.
- Ye Q, Wang B, Mao J. The pathogenesis and treatment of the 'Cytokine Storm' in COVID-19. *J Infect* 2020;80(6):607-13.
- Sauer A, Peukert K, Putensen C, Bode C. Antibiotics as immunomodulators: a potential pharmacologic approach for ARDS treatment. *Eur Respir Rev* 2021;30(162):210093.
- Feng LH, Li XD, Zhang XY, Cheng PJ, Feng ZY. Dexamethasone for the treatment of acute respiratory distress syndrome: A systematic review and meta-analysis. *Medicine (Baltimore)* 2022;101(39):e30195.
- Chen R, Lan Z, Ye J, Pang L, Liu Y, Wu W, et al. Cytokine storm: the primary determinant for the pathophysiological evolution of COVID-19 deterioration. *Front Immunol* 2021;12(2021):589095.
- Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir Med* 2020;8(4):420-2.
- Lingeswaran M, Goyal T, Ghosh R, Suri S, Mitra P, Misra S, et al. Inflammation, immunity and immunogenetics in COVID-19: a narrative review. *Indian J Clin Biochem* 2020;35(3):260-73.
- Widowati W, Darsono L, Suherman J, Fauziah N, Maesaroh M, Erawijantari PP. Anti-inflammatory effect of mangosteen (*Garcinia mangostana* L.) peel extract and its compounds in LPS-induced RAW264. 7 cells. *Nat Prod Sci* 2016;22(3):147-53.
- Sproston NR, Ashworth JJ. Role of C-reactive protein at sites of inflammation and infection. *Front Immunol* 2018;9(2018):754.
- Ohtsu A, Shibutani Y, Seno K, Iwata H, Kuwayama T, Shirasuna K. Advanced glycation end products and lipopolysaccharides stimulate interleukin-6 secretion via the RAGE/TLR4-NF- κ B-ROS pathways and resveratrol attenuate these inflammatory responses in mouse macrophages. *Exp Ther Med* 2017;14(5):4363-70.
- Boaru SG, Borkham-Kamphorst E, Van de Leur E, Lehnen E, Liedtke C, Weiskirchen R. NLRP3 inflammasome expression is driven by NF- κ B in cultured hepatocytes. *Biochem Biophys Res Commun* 2015;458(3):700-6.
- Danielski LG, Giustina A Della, Bonfante S, Barichello T, Petronilho F. The NLRP3 inflammasome and its role in sepsis development. *Inflammation* 2020;43(1):24-31.
- Pelaia C, Crimi C, Vatrella A, Tinello C, Terracciano R, Pelaia G. Molecular targets for biological therapies of severe asthma. *Front Immunol* 2020;11:603312.
- Carcattera M, Caruso C. Alveolar epithelial cell type II as main target of SARS-CoV-2 virus and COVID-19 development via NF-Kb pathway deregulation: A physiopathological theory. *Med Hypotheses* 2021;146:110412.
- Paudyal V, Sun S, Hussain R, Abutaleb MH, Hedima EW. Complementary and alternative medicines use in COVID-19: A global perspective on practice, policy and research. *Res Soc Adm Pharm* 2022;18(3):2524-8.
- Emaliyawati E, Nurhalimah H, Adha H, et al. Management of Acute Respiratory Distress Syndrome (ARDS) patients: A Literature Review. *Padjadjaran Acute Care Nurs J* 2021;2(2).
- Leyva-López N, Gutierrez-Grijalva EP, Ambriz-Perez DL, Heredia JB. Flavonoids as cytokine modulators: a possible therapy for inflammation-related diseases. *Int J Mol Sci* 2016;17(6):921.
- Novilla A, Djamhuri DS, Nurhayati B, Rihibiha DD, Afifah E, Widowati W. Anti-inflammatory properties of oolong tea (*Camellia sinensis*) ethanol extract and epigallocatechin gallate in LPS-induced RAW 264.7 cells. *Asian Pac J Trop Biomed* 2017;(12):1005-9.
- Souza NC, de Oliveira Nascimento EN, de Oliveira IB, et al. Anti-inflammatory and antioxidant properties of blend formulated with compounds of *Malpighia emarginata* DC (acerola) and *Camellia sinensis* L. (green tea) in lipopolysaccharide-stimulated RAW 264.7 macrophages. *Biomed Pharmacother* 2020;128:110277.
- Saeedi-Boroujeni A, Mahmoudian-Sani M-R. Anti-inflammatory potential of Quercetin in COVID-19 treatment. *J Inflamm (Lond)* 2021;18(1):3.
- Hidayat M, Prahastuti S, Afifah E, Widowati W, Yusuf M, Hasan K. The role of green peas protein hydrolysate in TGF/SMAD signaling to prevent renal fibrosis. *J King Saud Univ* 2022;34(4):101920.
- Girsang E, Ginting CN, Lister INE, Gunawan KY, Widowati W. Anti-inflammatory and antiaging properties of chlorogenic acid on UV-induced fibroblast cell. *PeerJ* 2021;9: e11419.
- An X, Sun X, Hou Y, Yang X, Chen H, Zhang P, et al. Protective effect of oxytocin on LPS-induced acute lung injury in mice. *Sci Rep* 2019;9(1):2836.
- Ngkelo A, Meja K, Yeadon M, Adcock I, Kirkham PA.

- LPS induced inflammatory responses in human peripheral blood mononuclear cells is mediated through NOX4 and G α dependent PI-3kinase signalling. *J Inflamm (Lond)* 2012;9(1):1.
25. Domscheit H, Hegeman MA, Carvalho N, Spieth PM. Molecular dynamics of lipopolysaccharide-induced lung injury in rodents. *Front Physiol* 2020;11:36.
 26. Huang Z, Kraus VB. Does lipopolysaccharide-mediated inflammation have a role in OA? *Nat Rev Rheumatol* 2016;12(2):123-9.
 27. Hawn TR, Verbon A, Janer M, Zhao LP, Beutler B, Aderem A. Toll-like receptor 4 polymorphisms are associated with resistance to Legionnaires' disease. *Proc Natl Acad Sci USA* 2005;102(7):2487-9.
 28. Ibrahim ZA, Armour CL, Phipps S, Sukkar MB. RAGE and TLRs: relatives, friends or neighbours? *Mol Immunol* 2013;56(4):739-44.
 29. Yu J, Bi X, Yu B, Chen D. Isoflavones: anti-inflammatory benefit and possible caveats. *Nutrients* 2016;8(6):361.
 30. Kim SM, Ha JS, Han AR, Cho S-W, Yang S-J. Effects of α -lipoic acid on LPS-induced neuroinflammation and NLRP3 inflammasome activation through the regulation of BV-2 microglial cells activation. *BMB Rep* 2019;52(10):613-8.
 31. Jo E-K, Kim JK, Shin D-M, Sasakawa C. Molecular mechanisms regulating NLRP3 inflammasome activation. *Cell Mol Immunol* 2016;13(2):148-59.
 32. Freeman TL, Swartz TH. Targeting the NLRP3 inflammasome in severe COVID-19. *Front Immunol* 2020;11:1518.
 33. Yan Z, Zhong Y, Duan Y, Chen Q, Li F. Antioxidant mechanism of tea polyphenols and its impact on health benefits. *Anim Nutr* 2020;6(2):115-23.
 34. Ohishi T, Goto S, Monira P, Isemura M, Nakamura Y. Anti-inflammatory action of green tea. *Antiinflamm Anti-Allergy Agents Med Chem (Formerly Curr Med Chem Anti-Allergy Agents)* 2016;15(2):74-90.
 35. Hodges JK, Sasaki GY, Bruno RS. Anti-inflammatory activities of green tea catechins along the gut-liver axis in nonalcoholic fatty liver disease: lessons learned from preclinical and human studies. *J Nutr Biochem* 2020:108478.
 36. He W, Qu T, Yu Q, Wang Z, Lv H, Zhang J, et al. LPS induces IL-8 expression through TLR 4, MyD88, NF-kappa B and MAPK pathways in human dental pulp stem cells. *Int Endod J* 2013;46(2):128-36.
 37. Noailles A, Maneu V, Campello L, Lax P, Cuenca N. Systemic inflammation induced by lipopolysaccharide aggravates inherited retinal dystrophy. *Cell Death Dis* 2018;9(3):350.
 38. Liao J, Xie X, Wang W, et al. Anti-inflammatory activity of essential oil from leaves of *Blumea balsamifera* (L.) DC through inhibiting TLR4/NF-kB signaling pathways and NLRP3 inflammasome activation in LPS-induced RAW264. 7 macrophage cells. *Journal of Essential Oil Bearing Plants* 2021;24(1):1-17.
 39. Kelley N, Jeltema D, Duan Y, He Y. The NLRP3 inflammasome: an overview of mechanisms of activation and regulation. *Int J Mol Sci* 2019;20(13):3328.
 40. Xue Y, Du M, Zhu M-J. Quercetin suppresses NLRP3 inflammasome activation in epithelial cells triggered by *Escherichia coli* O157:H7. *Free Radic Biol Med* 2017;108:760-9.
 41. Leu J-G, Lin C-Y, Jian J-H, Shih C-Y, Liang Y-J. Epigallocatechin-3-gallate combined with alpha lipoic acid attenuates high glucose-induced receptor for advanced glycation end products (RAGE) expression in human embryonic kidney cells. *An Acad Bras Cienc* 2013;85(2):745-52.
 42. Wang J, Fan SM, Zhang J. Epigallocatechin-3-gallate ameliorates lipopolysaccharide-induced acute lung injury by suppression of TLR4/NF-kB signaling activation. *Brazilian J Med Biol Res* 2019;52(7):e8092.
 43. Bao S, Cao Y, Zhou H, Sun X, Shan Z, Teng W. Epigallocatechin gallate (EGCG) suppresses lipopolysaccharide-induced Toll-like receptor 4 (TLR4) activity via 67 kDa laminin receptor (67LR) in 3T3-L1 adipocytes. *J Agric Food Chem* 2015;63(10):2811-9.
 44. Thao NP, Luyen BTT, Widowati W, Fauziah N, Maesaroh M, Herlina T, et al. Anti-inflammatory flavonoid C-Glycosides from *Piper aduncum* leaves. *Planta Med* 2016;82(17):1475-81.
 45. Patil KR, Mohapatra P, Patel HM, Goyal HM, Ojha SN, Kundu CN, et al. Pentacyclic triterpenoids inhibit IKK β mediated activation of NF-kB pathway: in silico and in vitro evidences. *PLoS One* 2015;10(5):e0125709.
 46. Paranga TG, Pavel-Tanasa M, Constantinescu D, Plesca CE, Petrovici C, Miftode I-L, et al. Comparison of C-reactive protein with distinct hyperinflammatory biomarkers in association with COVID-19 severity, mortality and SARS-CoV-2 variants. *Front Immunol* 2023;14:1213246.
 47. Salazar J, Martínez MS, Chávez-Castillo M, Núñez V, Añez R, Torres Y, et al. C-reactive protein: an in-depth look into structure, function, and regulation. *Int Sch Res Notices* 2014;2014:653045.
 48. Semple SJ. C-reactive protein—biological functions, cardiovascular disease and physical exercise. *South African Journal of Sports Medicine* 2006;18:24-8.
 49. Chen CH, Lin SW, Shen CF, Hsieh KS, Cheng CM. Biomarkers during COVID-19: mechanisms of change and implications for patient outcomes. *Diagnostics (Basel)* (2022) 12(2):509.
 50. Yang Y, Jiang G, Zhang P, Fan J. Programmed cell death and its role in inflammation. *Mil Med Res* 2015;2:12.
 51. Jiang K, Zhang T, Yin N, Ma X, Zhao G, Wu H, et al. Geraniol alleviates LPS-induced acute lung injury in mice via inhibiting inflammation and apoptosis. *Oncotarget* 2017;8(41):71038-53.
 52. Bagheri E, Hajiaghaalipour F, Nyamathulla S, Salehen N. The apoptotic effects of *Brucea javanica* fruit extract against HT29 cells associated with p53 upregulation and inhibition of NF-kB translocation. *Drug Des Devel Ther* 2018;12:657-71.

53. Qi D, Wang D, Zhang C, Tang X, He J, Zhao Y, et al. Vaspin protects against LPS-induced ARDS by inhibiting inflammation, apoptosis and reactive oxygen species generation in pulmonary endothelial cells via the Akt/GSK-3 β pathway. *Int J Mol Med* 2017;40(6):1803-17.
54. Tang J, Diao P, Shu X, Li L, Xiong L. Quercetin and quercitrin attenuates the inflammatory response and oxidative stress in LPS-induced RAW264. 7 cells: in vitro assessment and a theoretical model. *Biomed Res Int* 2019;2019.