

Development of an Immunoaffinity Method for Purification of Streptokinase

Zohreh Karimi ¹, Mohammad Babashamsi ^{2*}, Ezat Asgarani ¹, and Ali Salimi ²¹. Department of Biology, Faculty of Science, Alzahra University, Tehran, Iran². Department of Immunochemistry, Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

Abstract

Background: Streptokinase is a potent activator of plasminogen to plasmin, the enzyme that can solubilize the fibrin network in blood clots. Streptokinase is currently used in clinical medicine as a thrombolytic agent. It is naturally secreted by β -hemolytic *streptococci*.

Methods: To reach an efficient method of purification, an immunoaffinity chromatography method was developed that could purify the streptokinase in a single step with high yield. At the first stage, a CNBr-Ac-tivated sepharose 4B-Lysine column was made to purify the human blood plasminogen. The purified plasminogen was utilized to construct a column that could purify the streptokinase. The rabbit was immunized with the purified streptokinase and the anti-streptokinase (IgG) purified on another streptokinase substituted sepharose-4B column. The immunoaffinity column was developed by coupling the purified anti-Streptokinase (IgG) to sepharose 6MB-Protein A. The *Escherichia coli* (*E.coli*) BL21 (DE3) pLysS strain was transformed by the recombinant construct (cloned streptokinase gene in pGEX-4T-2 vector) and gene expression was induced by IPTG. The expressed protein was purified by immunoaffinity chromatography in a single step.

Results: The immunoaffinity column could purify the recombinant fusion GST-SK to homogeneity. The purity of streptokinase was confirmed by SDS-PAGE as a single band of about 71 kD and its biological activity determined in a specific streptokinase assay. The yield of the purification was about 94%.

Conclusion: This method of streptokinase purification is superior to the previous conventional methods.

* Corresponding author:

Mohammad Babashamsi,
Ph.D., Avicenna Research
Institute, ACECR, Tehran,
Iran

Tel: +98 21 22432020

Fax: +98 21 22432021

E-mail:

babashams@avicenna.ac.ir

Received: 1 Jan 2012

Accepted: 9 Apr 2012

Avicenna J Med Biotech 2012; 4(3): 142-147

Keywords: Chromatography, Purification, Streptokinase, Thrombolytic agent

Introduction

A failure of homeostasis and consequent formation of blood clots in circulatory system can produce severe outcomes such as stroke and myocardial infarction. Pathological development of blood clots requires clinical intervention with fibrinolytic agent such as urokinase, tissue plasminogen activator and streptokinase ⁽¹⁾.

Streptokinase (SK) is a non-protease plasminogen activator, produced by certain *streptococci* and certain bacteria which contain

appropriate genetic material derived from *streptococci* of Lancefield groups A, C or G. Tillet ⁽²⁾ discovered that this bacterial protein caused the lysis of human blood clots. The fibrinolytic activity of streptokinase originates in its ability to activate blood plasminogen to plasmin, the enzyme that degrades fibrin cloth through its specific lysine binding site ^(3,4). Due to this property it is used in clinical medicine as a therapeutic agent in the treatment of thromboembolic blockage, including

coronary thrombosis.

The mature streptokinase has a molecular weight of about 47 *kD* and was found to be composed of 415 amino acid residues⁽⁵⁾. Secretion of streptokinase into the external medium is directed by a 26 amino acid signal peptide which is cleaved during the secretion process.

Production of natural and recombinant forms of SK and its purification by different chromatography methods, which are based on quantitative differences in solubility, electrical charge, molecular size and shape or non specific physical interactions with surfaces, has been studied by several workers^(6,7). The growth of a β -hemolytic streptococcus was studied in continuous culture with pH as a limiting factor^(8,9).

We have already reported the rate of SK production in a fed-batch culture and purification by affinity chromatography on acylated plasminogen with *p*-nitro phenyl guanidine-benzoate (NPGB)⁽¹⁰⁾. The gene codes for streptokinase, from *Streptococcus equisimilis* (*S. equisimilis*) H46A was expressed in several heterologous gram positive and gram negative bacteria^(11,12). A fusion recombinant streptokinase has been produced and purified it in a single step affinity chromatography using glutathione as the ligand⁽¹³⁾.

In the present study, the issue of purification of recombinant Streptokinase (rSK) is addressed, for which a new immunoaffinity chromatography approach to produce active rSKC in a single step with high yield and stability was introduced.

Materials and Methods

The materials used in the experiment includes; H46A (ATCC 12449, USA), *E. coli* DH5 α and *E. coli* BL21 (DE3) pLysS (Invitrogen, USA), Todd Hewitt Broth (THB, HiMEDIA Laboratories), Trypticase Soy Agar (TSA, BBL, USA) Lysine monohydrochloride (Sigma Chemical, USA), pGEX-1.2-4T-2 (Avicenna Research Institute, Iran), Sodium Dodecyl Sulfate (SDS, Sigma Chemical, USA) Hexyl resorcinol (Merck, Germany),

NPGB (Sigma Chemical, USA), 3-amino-n-caproic acid (EACA, Sigma Chemical, USA), Chromogenic substrate (S-2251, Chromogenic laboratories, Italy), Isopropyl-beta-D-thiogalactopyranoside (IPTG, Roche, Germany), Cyanogen bromide-activated Sepharose 4B (Sigma Chemical, USA), Sepharose 6MB-Protein A (GE healthcare, USA), Dimethyl pimelimidate-HCL (DMP, sigma-Aldrich, USA), Phenylmethylsulfonyl fluoride (PMSF, sigma Chemical, USA), serum albumin (sigma-Aldrich, USA), sheep anti-rabbit IgG (Avicenna Research Institute, Iran), o-Phenylenediamine (OPD, sigma Chemical, USA), Polyvinylidene difluoride (PVDF, Roche, Germany), Enhanced Chemiluminescent substrates (ECL, GE Health care, Biotech Buckinghamshire, UK), Chemicals for PAGE (Sigma Chemical, USA), Salts for buffers (Merck, Germany).

Preparation of streptokinase-sepharose column

S. equisimilis group C, strain H46A was grown in TSA and THB. The secreted SK purified by protected affinity chromatography⁽¹⁰⁾. The purified streptokinase was then coupled to CNBr-activated Sepharose 4B. Briefly, 5 mg of purified SK was added per ml of the CNBr-activated Sepharose 4B resin in 0.1 M Sodium bicarbonate (NaHCO₃, pH=8.3) and after 2 hr at room temperature, the resin was washed with 0.2 M glycine, (pH=8) and then alternately for three times with 0.1 M sodium acetate, 0.5 M NaCl, (pH=4) and 0.1 M NaHCO₃, (pH=8.3). The coupled streptokinase-sepharose was transferred to column, neutralized by Phosphate Buffered Saline (PBS) and stored in PBS-azide.

Production and purification of anti-streptokinase

Rabbit was immunized by subcutaneous injection of 1500 μ l of an emulsion containing 25 μ g of streptokinase and equal volume of complete Freund's adjuvant at 3 weeks intervals. Blood collected and serum was separated. Ten ml of sera was diluted 5 times with PBS, centrifuged at 4000 rpm for 10 min at 4°C to remove residual particles, and then was filtered. The supernatant was passed through a

SK-Sepharose column at a rate of 12 ml per hour. It was washed by PBS and eluted with 0.2 M Glycine- HCl pH=2.5. Then it was neutralized by PBS. The purity was confirmed by SDS-PAGE.

Expression tests of fusion GST-SK in *E.coli*

SDS-PAGE analysis: *E.coli* BL21 (DE3) pLysS was transformed by pGEX-1.2-4T-2 (pGEX vector with Glutathione S-Transferase (GST) tag and SK gene insert) ⁽¹³⁾. Bacteria was grown in 5 mL LB broth to an OD of 0.6 (A600 nm) with vigorous agitation at 37°C. Fusion protein expression was induced by 0.1 mM IPTG and incubated for an additional 7 hr. One ml of the culture was collected at two hours intervals (1, 3, 5 and 7 hr), spun down and pellet was resuspended in 100 µl of 1x SDS sample buffer containing (7.5% Tris. base, 2 ml of 10% SDS, 1 ml Glycerol, 2 mg Bromophenol blue, 25 µl 2Mercaptoethanol). The cells were lysed using a strong vortex, heated at 100°C for 5 min, spun down and its supernatant was applied to 10% SDS- polyacrylamide gel electrophoresis. Proteins were stained with Coomassie blue and the band was visualized by destaining. In order to analyse the fusion GST-SK protein Molecular Weight (MW), the Bio-Rad MW marker, Cat. No. 161-0309 was used.

Western blotting: Sonicates of pre and post inductions of cloned GST-SK in *E.coli* and also a purified SK were run on SDS-PAGE. The gel was blotted on PVDF membrane using transfer buffer containing 25 mM Tris (pH=8.3), 192 mM glycine and 20% methanol at 100 v for 75 min at room temperature. Blotted membrane was blocked with 2.5% skim milk in TBST buffer (0.5 M NaCl, 0.02 M Tris pH=8.5, 0.05% tween 20) overnight at 4°C. After twice washing with PBS-Tween, membranes were incubated for 1.5 hr at room temperature with 1 µg per ml rabbit antistreptokinase in 1% skim milk. After reactions with primary antibody, the membrane was washed three times with PBS-Tween and incubated with peroxidase conjugated sheep anti-rabbit IgG (Avicenna Research Institute)

at a 1:2000 dilution in 1% skim milk. The membrane was then washed three times with PBS-Tween. The reaction was developed by ECL for 1 min and scanned.

Preparation of antistreptokinase-sepharose column

Cross-linking of antistreptokinase to protein-A sepharose 6MB: Twenty mg of purified anti-SK antibody in 10 ml of PBS was mixed with 1 ml of protein A-Sepharose 6MB continuously at 4°C and washed twice with excess of PBS. The beads rinsed with 8 ml of 0.2 M borate-NaOH, pH=8.6 and centrifuged for 1 min at 500×g. The washing was repeated with borate buffer two more times. Anti-SK was cross-linked to protein-A Sepharose by addition of 15.5 mg of DMP in 2 ml of 0.2 M triethanolamin buffer, pH=8.3. to 1 ml of the coupled bead. After rotation for 30 min at room temperature, the cross-linking and washing steps were repeated twice to improve cross-linking efficiency. The rest of reactive amino groups were quenched by addition of 50 mmol/l glycine-HCL (pH=3). The antibodies cross-linked to beads were stored at 4°C in PBS containing 0.02% sodium-Azide Tween-20 (PBST) until use.

Purification of streptokinase on antistreptokinase-sepharose column: Two ml of 50% slurry of antibody-cross linked to protein A-sepharose was equilibrated with 20 ml of binding buffer [10 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES)-NaOH, pH=7, 0.1 M NaCl, 2 mM PMSF]. Ten ml of recombinant streptokinase sample [sonicate of a 6 hours post induction of transformed BL21 (DE3) pLysS] was prepared in binding buffer and added on the bead. The binding occurred for 1 hr to over-night by continuously mixing. The bead was packed in column and sequentially washed with 30 ml of TBS (20 mM Tris-HCl, pH=7.5. 150 mM NaCl) with a flow rate of 1 ml/min and then with 10 ml pre-elution buffer (0.1 X TBS). The bound proteins were eluted with 5 ml of 50 mM sodium phosphate, pH=12. Five 1 ml fractions collected into tubes preloaded with 1 ml of cold

1 M Tris-HCl, pH=6. As soon as each fraction volume reaches 2 ml, contents of each tube were mixed to neutralize the eluate and placed on ice. The column was regenerated immediately after use by washing with 40 ml of 1 M Tris-HCl.

The activity of purified SK was measured by plasmin hydrolysis of chromogenic peptidyl anilide substrate (S-2251), procedure⁽¹⁴⁾ with slight modification by us⁽¹³⁾.

Results

The culture of BL21 (DE3) PlysS', transformed by PGEX-1.2-4T2 construct induced by IPTG indicates that, the 71 kD fusion GST-SK protein is expressed increasingly (Figure 1). The western blot analysis of pre and post induction of this bacterial culture approved that the expressed protein is GST-SK (Figure 2).




Figure 1. 10% SDS-PAGE; fusion GST-SK expression in *E. coli*, BL21 PLYsS: Lane 1: Pre-induction. Lane 3: 2 hr, Lane 4: 4 hr, Lane 5: 6 hr and Lane 6: 8 hr post-induction. Lane 7: Protein marker (Fermentas SM0441). Lane 8: Pure SK




Figure 2. Western Blot of fusion GST-SK expression in *E. coli*, BL21 PLYsS: Lane 1: Protein MW marker (Fermentas SM0671). Lanes 2, 3, 4, 5 and 6: Cloned GST-SK in *E. coli*, after induction. Lane 7: BSA. Lanes 8, 9 and 10: Pure SK

The SK-Sepharose column constructed and the anti-SK antibody produced in rabbit and purified on the SK-Sepharose column. The purified antibody loaded on SDS-PAGE and the purity was confirmed (Figure 3).

The SK produced by H46a was purified by immunoaffinity column and two distinct bands of SK isotypes (48 and 45 kD) observed on SDS-PAGE (Figure 4). The im-




Figure 3. 10% SDS-PAGE; purification of anti-SK from rabbit sera, Lane 1: BSA, Lane 2: Flow-through, Lane 3: serum, Lanes 4, 6, and 7: Pure anti-SK, Lane 8: IgG




Figure 4. 10% SDS-PAGE; purification of SK from H46a, by immunoaffinity column Lane 1: Cultured H46a, Lane 2: Flow-through, Lane 3: Protein marker, Lane 4 and 5: Purified SK, Lane 6: BSA




Figure 5. 10% SDS-PAGE; purification of recombinant fusion GST-SK from *E. coli*, by immunoaffinity column. Lane 1: Purified GST-SK 73 kD, Lane 2: Flow-through, Lane 3: Sonicate of *E. coli*, Lane 4: Protein marker (Fermentas SM0441), Lane 5: Hand made marker (Reduced IgG and BSA), Lane 6: BSA, Lanes 7 and 8: Non-transformed *E. coli*

Table 1. Purification of fusion GST-SK

Steps	Total recovery		Specific activity	Fold purified	% Yield
	Protein (mg)	SK (IU)			
Initial (Sonicate)	12.7	120500	9488	--	--
Elution (Purified)	4.25	113270	26652	2.8	94

mune affinity column could purify the recombinant Fusion GST-SK to homogeneity as a single band of about 71 kD on SDS-PAGE (Figure 5). The yield of the purification was about 94% (Table 1).

Discussion

Purification of SK have already been studied by various methods including; column chromatography on DEAE-cellulose followed by column electrophoresis in sucrose density gradients ⁽¹⁵⁾, a combination of ion exchange (DEAE-Sephadex A-50) and gel permeation (Sephadex G-100) chromatography ⁽¹⁶⁾. These methods required repeated chromatography to remove the impurities. As a result, the SK recovery yields found to be 40-60%.

In other attempts SK was fractionated by ammonium sulfate and further purified by gradient elution from a DEAE-cellulose chromatography column ⁽¹⁷⁾. Several affinity chromatography methods were applied by different workers, including; insolubilized diisopropyl fluorophosphates (DIP) plasmin as the affinity ligand ⁽¹⁸⁾. This method caused a 30% decrease in the streptokinase activity.

As mentioned earlier, SK was purified by a protected affinity chromatography method in which the ligand (plasmin) was acylated to protect its activation by SK ⁽¹⁰⁾. Even though the yield of purification by this method was found to be 95%, but acylation is not permanent and remains stable just for less than 4 hours. We have produced a recombinant SK tagged by a poly Histidin tail and purified both of the 6xHis-SK and GST-SK by affinity chromatography on NiNTA-agarose and glutathione-sepharose columns respectively, using thrombin for cleavage of the tags (unpublished). Thrombin is a costly enzyme, so

this method is not cost effective for large scale purification.

Recently, we have extracted and purified the native SK from H46a fermentation broth by chemical reduction method based on the specific structure of SK, lacking disulfide bridge ⁽¹⁹⁾. This method may not be applicable for recombinant SK purification from other bacterial sources, as they may secrete proteins lacking disulfide bridges.

Conclusion

Most of the mentioned methods have limitations, either they are multisteps and often results in unacceptable losses of SK or inadequate removal of impurities. The present immunoaffinity method based on reversible linkage among antigen and antibody could purify the natural streptococcal SK and recombinant GST-SK in a single step with a yield of 94%. This method of SK purification is superior to the previous conventional methods as it does not require the addition of a tag to the recombinant SK for affinity purifications. Moreover, this method does not require any chemical modification.

In usual immunoaffinity method for purification of an antigen, the column consists of the immobilized antibody to the resin through its Fab region. So, the number of free Fab domains for antigen-antibody interaction is limited.

The present technique utilized protein A-sepharose 6MB as solid support to increase the chance of SK interaction with the Fab region of anti-SK, so antibody adhered to protein A from Fc region. A specific linker protected the anti-SK cleavage by elution buffer.

Acknowledgement

We would like to thank research deputy of

the Academic Center for Education, Culture and Research (ACECR) for financial support (Grant No: 1072-21).

References

- Banerjee A, Chistic Y, Banerjee UC. Streptokinase, a clinically useful thrombolytic agent. *Biotechnol Adv* 2004;22(4):287-307.
- Tillet WS, Garner RL. The fibrinolytic activity of hemolytic streptococci. *J Exp Med* 1933;58(4):485-502.
- Castellino FJ. A unique enzyme-protein substrate modifier reaction: plasmin/streptokinase interaction. *Trends Biochem Sci* 1979;4(1):1-5.
- Rodriguez P, Fuentes P, Barro M, Alvarez JG, Muñoz E, Collen D, et al. Structural domains of streptokinase involved in the interaction with plasminogen. *Eur J Biochem* 1995;229(1):83-90.
- Jackson KW, Tang J. Complete amino acid sequence of streptokinase and its homology with serine proteases. *Biochemistry* 1982;21(26):6620-6625.
- Blatt WF, Segal H, Gray JL. Purification of streptokinase and human plasmin and their interaction. *Thromb Diath Haemorrh* 1964;11:393-403.
- Jeong IK, Young EL, Jeung SJ, Myung B. Purification of streptokinase by affinity chromatography using human plasminogen. *Korean Biochem J* 1993;26(2):172-175.
- Karush F, Lacocca VF, Harris TN. Growth of group A hemolytic streptococcus in the steady state. *J Bacteriol* 1956;72(3):283-294.
- Ogburn CA, Harris TN, Harris S. Extracellular antigens in steady-state cultures of the hemolytic streptococcus: production of proteinase at low pH. *J Bacteriol* 1958;76(2):142-151.
- Babashamsi M, Razavian MH, Nejadmoghadam MR. Production and purification of streptokinase by protected affinity chromatography. *Avicenna J Med Biotech* 2009;1(1):47-51.
- Steiner K, Malke H. Dual control of streptokinase and streptolysin S production by the covRS and fasCAX two-component regulators in *Streptococcus dysgalactiae* subsp. *equisimilis*. *Infect Immun* 2002;70(7):3627-3636.
- Kaur J, Rajamohan G, Dikshit KL. Cloning and characterization of promoter-active DNA sequences from *Streptococcus equisimilis*. *Curr Microb* 2007;54(1):48-53.
- Nejadmoghadam MR, Modarresi MH, Babashamsi M, Chamankhah M. Cloning and over expression of active recombinant fusion streptokinase: a new approach to facilitate purification. *Pak J Biol Sci* 2007;10(13):2146-2151.
- Jackson KW, Esmon N, Tang J. Streptokinase and staphylokinase. *Methods Enzymol* 1981;80:387-394.
- De Renzo EC, Suteri PK, Hutchings BL, Bell PH. Preparation and certain properties of highly purified streptokinase. *J Biol Chem* 1967;242(3):533-542.
- Taylor FB, Botts J. Purification and characterization of streptokinase with studies of streptokinase activation of plasminogen. *Biochemistry* 1968;7:232-242.
- Einarsson M, Skoog B, Forsberg B, Einarsson R. Characterization of highly purified native streptokinase and altered streptokinase after alkaline treatment. *Biochimica et Biophysica Acta* 1979;568(1):19-29.
- Castellino FJ, De Renzo EC, Suteri PK, Hutchings BL, Bell PH. Preparation and certain properties of highly purified streptokinase. *J Biol Chem* 1967;242:533-542.
- Karimi Z, Babashamsi M, Asgarani E, Niakan M, Salimi A. Fermentation, fractionation and purification of streptokinase by chemical reduction method. *Iran J Microbiol* 2011;3(1):42-46.